

12.804 — One Layer Inversion/ Rossby Waves — Num. Expts.

Model

$$\frac{\partial}{\partial t}q' + J(\psi, q') + \beta \frac{\partial}{\partial x}\psi = \textit{filtering} \quad (1)$$

$$q' = \nabla^2\psi - F_1\psi \quad (2)$$

The filtering is required for numerical stability and is intended to remove small scale enstrophy without dissipating too much energy — a typical numerical trick. The parameter F_1 is zero for the barotropic vorticity equation and non-zero for the shallow water QG system. The model works on a doubly periodic domain with $-\pi \leq x < \pi$, $-\pi \leq y < \pi$. It is a pseudospectral model, meaning the inversion of (2) is done by working in Fourier space, as is the evaluation of the frictional term, the β term, and the time-stepping. The advection term is rewritten as

$$\nabla \cdot (\vec{u}q')$$

and is solved by evaluating the transform of the velocities [multiplying the transform of ψ by $(-i\ell, ik)$], transforming them to real space and multiplying by the real space value of q' . The result is transformed back to wavenumber space and dotted with $(ik, i\ell)$.

PV inversion

The interface allows you to specify the q' values and determine the resulting ψ fields. For the inversion, you specify the parameters F_1 and β . You express the PV anomaly q' as functions of the x and y matrices and then the program will calculate ψ and contour both the PV anomaly q' and the full PV field $q' + \beta y$. For example, consider

$$q=3*\cos(2*x-3*y)+0.01*\sin(x)$$

The program can also be run in the opposite direction: specifying ψ and calculating q , e.g.

$$\text{psi}=1.5*\exp(-2*x.^2-2*y.^2);$$

In addition you can give values to the parameters β and F_1 ($= r_d^{-2}$).

PV evolution

Once you have specified the PV and/or streamfunction field, use the link appropriate to your machine to see how the flow evolves. You can alter the time step $\Delta t = 1/nsteps$, the contour interval ci for the display, and the total time length for the run $timelength$. The contours will be shown every $nsnap$ time steps.

When you're finished be sure to hit the Quit button before closing the browser.

Things you can try

- 1) Verify the dispersion relation with and without F_1 . Look at how waves with their crests at an angle propagate. Remember — initial conditions should be periodic in x and y with period 2π .
- 2) Consider how superpositions of two waves propagate. When does nonlinearity enter?
- 3) Look at the dispersion of one- and two-dimensional isolated features. How can you understand the latter? How does nonlinearity affect it?
- 4) What happens to an elliptical eddy on an f -plane? A β -plane?

```
psi=exp(-2*x.^2-4*y.^2);  
f1=0;  
beta=0 to 0.5;
```

- 5) Look at the stability of a Rossby wave

```
psi=0.2*cos(2*x-3*y)+0.01*sin(x);  
timelength=30;  
f1=0.0;  
beta=1;
```

- 6) Look at the stability of a vortex
- 7) Examine the propagation of dipoles with and without β .

MIT OpenCourseWare
<http://ocw.mit.edu>

12.804 Large-scale Flow Dynamics Lab
Fall 2009

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.