

Recitation 10
October 12, 2010

- **Question 1.** The two parts of this question are about identities for a probabilistic model with sample space Ω , events A and B , and discrete random variable X . Any time conditioning on an event is indicated, the event has positive probability. An identity is *true* when it holds without any additional restrictions; it is *false* when there is any counterexample.

1.1. Which **one** of the following statements is **true**?

- (a) $\mathbf{P}(A \cap B)$ may be larger than $\mathbf{P}(A)$.
- (b) The variance of X may be larger than the variance of $2X$.
- (c) If $A^c \cap B^c = \emptyset$, then $\mathbf{P}(A \cup B) = 1$.
- (d) If $A^c \cap B^c = \emptyset$, then $\mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B)$.
- (e) If $\mathbf{P}(A) > 1/2$ and $\mathbf{P}(B) > 1/2$, then $\mathbf{P}(A \cup B) = 1$.

1.2. Which **one** of the following statements is **true**?

- (a) If $\mathbf{E}[X] = 0$, then $\mathbf{P}(X > 0) = \mathbf{P}(X < 0)$.
- (b) $\mathbf{P}(A) = \mathbf{P}(A | B) + \mathbf{P}(A | B^c)$
- (c) $\mathbf{P}(B | A) + \mathbf{P}(B | A^c) = 1$
- (d) $\mathbf{P}(B | A) + \mathbf{P}(B^c | A^c) = 1$
- (e) $\mathbf{P}(B | A) + \mathbf{P}(B^c | A) = 1$

- **Question 2.** Provide **clear reasoning**; partial credit is possible

Heather and Taylor play a game using independent tosses of an unfair coin. A head comes up on any toss with probability p , where $0 < p < 1$. The coin is tossed repeatedly until either the second time head comes up, in which case Heather wins; or the second time tail comes up, in which case Taylor wins. Note that a full game involves 2 or 3 tosses.

- 2.1. Consider a probabilistic model for the game in which the outcomes are the sequences of heads and tails in a full game. Provide a list of the outcomes and their probabilities of occurring.
- 2.2. What is the probability that Heather wins the game?
- 2.3. What is the conditional probability that Heather wins the game given that head comes up on the first toss?
- 2.4. What is the conditional probability that head comes up on the first toss given that Heather wins the game?

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
Department of Electrical Engineering & Computer Science
6.041/6.431: Probabilistic Systems Analysis
(Fall 2010)

- **Question 3.** Provide **clear reasoning**; partial credit is possible

A casino game using a **fair** 4-sided die (with labels 1, 2, 3, and 4) is offered in which a **basic game** has 1 or 2 die rolls:

- If the first roll is a 1, 2, or 3, the player wins the amount of the die roll, in dollars, and the game is over.
- If the first roll is a 4, the player wins \$2 and the amount of a second (“bonus”) die roll in dollars.

Let X be the payoff in dollars of the basic game.

3.1. Find the PMF of X , $p_X(x)$.

3.2. Find $\mathbf{E}[X]$.

3.3. Find the conditional PMF of the result of the first die roll given that $X = 3$. (Use a reasonable notation that you define explicitly.)

3.4. Now consider an **extended game** that can have any number of bonus rolls. Specifically:

- * Any roll of a 1, 2, or 3 results in the player winning the amount of the die roll, in dollars, and the termination of the game.
- * Any roll of a 4 results in the player winning \$2 and continuation of the game.

Let Y denote the payoff in dollars of the extended game. Find $\mathbf{E}[Y]$.

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.