

6.041SC Probabilistic Systems Analysis and Applied Probability, Fall 2013 Transcript – Recitation: Markov Chain Practice 1

Hi, everyone. Today, I'm going to talk about Markov Chain Practice number one. Before we start, let's first take a look at this Markov chain. This Markov chain has six states. In this problem, we always assume the process starts from state S_0 . On the first trial, the process can either make a transition from S_0 to S_1 with probability $1/3$ or from S_0 to S_3 with probability $1/3$ or from S_0 to S_5 with probability $1/3$.

If on the first trial, the process makes the transition from S_0 to S_1 or from S_0 to S_5 , it will always be stuck in either S_1 or S_5 forever, because both of the states S_1 and S_5 have a self-transition probability of one. On the other hand, if on the first trial, the process makes the transition from S_0 to S_3 , it can then either transition to the left or transition to the right or make self-transition back to the state S_3 .

If the process ever enters the left of the chain, it will never be able to come to the right. On the other hand, if the process ever enters the right of the chain, it would never be able to go to the left. For part A of the problem, we have to calculate the probability that the process enters S_2 for the first time at the case trial.

First, notice that it would take at least two trials for the process to make a transition from S_0 to S_2 . Therefore, for k equal to 1, the probability of a_k is simply equal to 0. For k equal to 1, probability of a_1 is equal to 0. Then for k equal to 2, 3 and on, the probability that the process enters S_2 for the first time at a case trial is equivalent to the probability that the process first makes a transition from S_0 to S_3 and then stays in S_3 for the next two k minus 2 trials and finally makes a transition from S_3 to S_2 on the k th trial.

So let's write this out. For k equal to 2, 3, and on, the probability of a_k is equal to the probability that the process first makes transition from S_0 to S_3 on the first trial, which is probability $1/3$, times the probability that the process makes self-transition for the next k minus 2 trials, which is probability $1/4$ to the power of k minus 2, and finally makes a transition from S_3 to S_2 on the k th trial, which is $1/4$. And this gives us $1/3$ times $1/4$ to the power of k minus 2 times $1/4$, which is equal to $1/3$ times $1/4$ to the power of k minus--

For part B of the problem, we have to calculate the probability that the process never enters as four. This event can happen in three ways. The first way is that the process makes a transition from S_0 to S_1 on the first trial and be stuck in S_1 forever. The second way that the process makes a transition from S_0 to S_5 on the first trial and be stuck at S_5 forever. The third way is that the process makes a transition from S_0 to S_3 on the first trial and then it makes a transition from S_3 to S_2 on the next state change so that it would never be able to go to S_4 .

Therefore, the probability of B is equal to the sum of probabilities of this three events. So the probability of B is equal to the probability that the process makes a transition from S_0 to S_1 on the first trial, which is $1/3$, plus the probability that the process makes a transition from S_0 to S_5

on the first trial, which is also $1/3$, plus the probability that the process makes a transition from S_0 to S_3 on the first trial times the probability that the process then makes a transition from S_3 to S_2 on the next state change.

So transition to S_2 , given that the processes are already in state S_3 and there's a state change. Let's take a look at this conditional probability. The condition that the processes are already in state S_3 and there's a state change imply two possible events, which are the transition from S_3 to S_2 and the transition from S_3 to S_4 . Therefore, we can write this conditional probability as the conditional probability of transition from S_3 to S_2 , given that another event, S_3 to S_2 or S_3 to S_4 has happened.

And this is simply equal to the proportion of p_{32} and p_{32} plus p_{34} , which is equal to $1/4$ over $1/4$ plus $1/2$, which is equal to $1/3$. Therefore, the probability of B is equal to $1/3$ plus $1/3$ plus $1/3$ times the $1/3$ here, which is equal to $7/9$. For part C of the problem, we have to calculate the probability that the process enter S_2 and leaves S_2 on the next trial.

This probability can be written as the product of two probabilities-- the probability that the process enters S_2 and the probability that it leaves S_2 on the next trial, given it's already in S_2 . Let's first look at the probability that the process enters S_2 . Using a similar approach as part B, we know that the probability the process ever enters S_2 is equal to the probability of the event that the process first makes a transition from S_0 to S_3 on the first trial and then makes a transition from S_3 to S_2 on the next state change.

So the probability that the process enters S_2 is equal to the probability that it first makes a transition from S_0 to S_3 on the first trial, which is P_{03} , times the probability that it makes a transition to S_2 , given that it's already in S_3 and there is a state change. We have already calculated this conditional probability in part B. Let's then look at the second probability term, the probability that the process leaves S_2 on the next trial, given that it's already in S_2 .

So given that the process is already in S_2 , it can take two transitions. It can either transition from S_2 to S_1 or make a self-transition from S_2 back to S_2 . Therefore, this conditional probability that it leaves S_2 on the next trial, given that it was already in S_2 is simply equal to the transition probability from S_2 to S_1 , which is P_{21} . Therefore, this is equal to P_{03} , which is $1/3$, times $1/3$ from the result from part B times P_{21} , which is $1/2$, and gives us $1/18$.

For part D of the problem, we have to calculate the probability that the process enters S_1 for the first time on the third trial. So if you take a look at this Markov chain, you'll notice that the only way for this event to happen is when a process first makes a transition from S_0 to S_3 on the first trial and from S_3 to S_2 on the second trial and from S_2 to S_1 on the third trial. Therefore, the probability of D is equal to the probability of the event that the process makes a transition from S_0 to S_3 on the first trial and from S_3 to S_2 on the second trial and finally from S_2 to S_1 on the third trial.

So this is equal to P_{03} times P_{32} times P_{21} , which is equal to $1/3$ times $1/4$ times $1/2$, which is equal to $1/24$. For part E of the problem, we have to calculate the probability that the process is in S_3 immediately after the n th trial. If you take a look at this Markov chain, you'll notice that if

on the first trial, the process makes a transition from S_0 to S_1 or from S_0 to S_5 , it will never be able to go to S_3 .

On the other hand, if on the first trial, the process makes a transition from S_0 to S_3 and if it leaves S_3 at some point, it will never be able to come back to S_3 . Therefore, in order for the process to be S_3 immediately after the n th trial, we will need the process to first make transition from S_0 to S_3 on the first trial and then stay in S_3 for the next n minus 1 trials. Therefore, the probability of the event e is simply equal to the probability of this event, which is equal to P_{03} times P_{33} to the power n minus 1, which is equal to $1/3$ times $1/4$ to the power of n minus 1.

And this concludes our practice on Markov chain today.

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.