

6.641 Electromagnetic Fields, Forces, and Motion  
Spring 2009

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.


Problem Set #3  
 Spring Term 2009

Issued: 2/19/09  
 Due: 2/26/09

Video Viewing: Haus/Melcher/Zahn Demos 8.4.1, and 8.6.1

Problem 3.1

An electric dipole consists of two opposite polarity charges,  $\pm q$  at  $z = \pm d/2$ .



- (a) Start with the electric potential of a point charge, and determine  $\Phi(r, \theta)$  for the electric dipole.
- (b) Define the dipole moment as  $p=qd$  and show that in the limit where  $d \rightarrow 0$  (while  $p$  remains finite), the electric potential is

$$\Phi(r, \theta) = \frac{p}{4\pi\epsilon_0} \frac{\cos \theta}{r^2}$$

- (c) What is the electric field for the dipole of part (b) with  $d \rightarrow 0$  with  $p$  remaining finite?
- (d) The electric field lines are lines that are tangent to the electric field:

$$\frac{dr}{rd\theta} = \frac{E_r}{E_\theta}$$

Using the result of (c), integrate this equation to find the field line that passes through the radial point  $r_0$  when  $\theta = \pi/2$ . This analytical equation can be used to precisely plot the electric field lines.

Hint:  $\int \cot \theta d\theta = \ln(\sin \theta) + \text{constant}$

- (e) Use your favorite computer plotting routine to plot on the same plot the equipotential and electric field lines for  $4\pi\epsilon_0 / p = 100 \text{ volt}^{-1}\cdot\text{m}^{-2}$ . Draw electric field lines for  $r_0 = 0.25, 0.5, 1$  and  $2$  meters and draw equipotential lines for  $\Phi = 0, \pm .0025, \pm .01, \pm .04, \pm .16$  and  $\pm .64$  volts.

### Problem 3.2


When a bird perches on a dc high-voltage power line and then flies away, it does so carrying a net charge.

- Why?
- For the purpose of measuring this net charge  $Q$  carried by the bird, we have the apparatus pictured below. Flush with the ground, a strip electrode having width  $w$  and length  $l$  is mounted so that it is insulated from ground. The resistance,  $R$ , connecting the electrode to ground is small enough that the potential of the electrode (like that of the surrounding ground) can be approximated as zero. The bird flies in the  $x$  direction at a height  $h$  above the ground with a velocity  $U$ . Thus, its position is taken as  $y=h$  and  $x=Ut$ . At time  $t$ , what is the effective charge distribution that will allow easy calculation of the electric scalar potential?
- The bird flies at an altitude  $h$  sufficiently large to make it appear as a point charge. What is the potential distribution as a function of time and position  $(x, y, z)$ ?
- Determine the surface charge density  $\sigma_s(x, y=0, z, t)$  on the ground plane at  $y=0$  as a function of time.
- At time  $t$ , what is the net charge,  $q$ , on the electrode? (Assume that the width  $w$  is small compared to  $h$  so that in an integration over the electrode surface, the integration in the  $z$  direction is simply a multiplication by  $w$ .)

Hint: Let  $x' = x - Ut$

$$\text{Hint: } \int \frac{dx}{[a^2 + x^2]^{3/2}} = \frac{x}{a^2[a^2 + x^2]^{1/2}}$$


- The current through the resistor is  $dq/dt$ . Find an expression for the voltage,  $v$ , that would be measured across the resistance,  $R$ .



"Bird on Powerline" diagram from: *Electromagnetic Fields and Energy* by Hermann A. Haus and James R. Melcher.


Problem 3.3

Find the magnetic field intensity at the point  $P$  shown due to the following line currents:




Problem 3.4

A constant current  $K_0 \bar{i}_\phi$  flows on the surface of a sphere of radius  $R$ .



- What is the magnetic field intensity at the center of the sphere?  
(Hint:  $\bar{i}_\phi \times \bar{i}_r = \cos\theta \cos\phi \bar{i}_x + \cos\theta \sin\phi \bar{i}_y - \sin\theta \bar{i}_z$ )
- Use the results of (a) to find the magnetic field intensity at the center of a spherical shell of inner radius  $R_1$  and outer radius  $R_2$  carrying a uniformly distributed volume current  $\bar{J} = J_0 \bar{i}_\phi$ .

Problem 3.5



A line current  $I$  of infinite extent in the  $z$ -direction is at a distance  $d$  above a perfectly conducting plane.

- Use the method of images to satisfy boundary conditions and find the magnetic vector potential for  $y > 0$ .
- What is the magnetic field for  $y > 0$ ?
- What is the surface current distribution that flows on the  $y=0$  surface?
- What is the force per unit length on the line current at  $y=d$ ?