18.06 Final Exam May 18, 2010 Professor Strang Your PRINTED name is: 1. Your recitation number is 2. 3. 4. 5. 6. 7. 8. 9. 9.

1. (12 points) This question is about the matrix

$$A = \left[\begin{array}{rrrr} 1 & 2 & 0 & 1 \\ 2 & 4 & 1 & 4 \\ 3 & 6 & 3 & 9 \end{array} \right].$$

- (a) Find a lower triangular L and an upper triangular U so that A = LU.
- (b) Find the reduced row echelon form R = rref(A). How many independent columns in A?
- (c) Find a basis for the nullspace of A.
- (d) If the vector b is the sum of the four columns of A, write down the complete solution to Ax = b.

- 2. (11 points) This problem finds the curve $y = C + D 2^t$ which gives the best least squares fit to the points (t, y) = (0, 6), (1, 4), (2, 0).
 - (a) Write down the 3 equations that would be satisfied *if* the curve went through all 3 points.

(b) Find the coefficients C and D of the best curve $y = C + D2^t$.

(c) What values should y have at times t = 0, 1, 2 so that the best curve is y = 0?

- 3. (11 points) Suppose $Av_i = b_i$ for the vectors v_1, \ldots, v_n and b_1, \ldots, b_n in \mathbb{R}^n . Put the v's into the columns of V and put the b's into the columns of B.
 - (a) Write those equations $Av_i = b_i$ in matrix form. What condition on which vectors allows A to be determined uniquely? Assuming this condition, find A from V and B.

(b) Describe the column space of that matrix A in terms of the given vectors.

(c) What additional condition on which vectors makes A an *invertible* matrix? Assuming this, find A^{-1} from V and B.

4. (11 points)

(a) Suppose x_k is the fraction of MIT students who prefer calculus to linear algebra at year k. The remaining fraction $y_k = 1 - x_k$ prefers linear algebra.

At year k + 1, 1/5 of those who prefer calculus change their mind (possibly after taking 18.03). Also at year k + 1, 1/10 of those who prefer linear algebra change their mind (possibly because of this exam).

Create the matrix A to give
$$\begin{bmatrix} x_{k+1} \\ y_{k+1} \end{bmatrix} = A \begin{bmatrix} x_k \\ y_k \end{bmatrix}$$
 and find the limit of $A^k \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ as $k \to \infty$.

(b) Solve these differential equations, starting from x(0) = 1, y(0) = 0:

$$\frac{dx}{dt} = 3x - 4y \quad \frac{dy}{dt} = 2x - 3y.$$

(c) For what initial conditions $\begin{bmatrix} x(0) \\ y(0) \end{bmatrix}$ does the solution $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$ to this differential equation lie on a single straight line in R^2 for all t?

5. (11 points)

- (a) Consider a 120° rotation around the axis x = y = z. Show that the vector i = (1,0,0) is rotated to the vector j = (0,1,0). (Similarly j is rotated to k = (0,0,1) and k is rotated to i.) How is j i related to the vector (1,1,1) along the axis?
- (b) Find the matrix A that produces this rotation (so Av is the rotation of v). Explain why $A^3 = I$. What are the eigenvalues of A?
- (c) If a 3 by 3 matrix P projects every vector onto the plane x+2y+z=0, find three eigenvalues and three independent eigenvectors of P. No need to compute P.

6. (11 points) This problem is about the matrix

$$A = \left[\begin{array}{rrr} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{array} \right].$$

- (a) Find the eigenvalues of $A^T A$ and also of $A A^T$. For both matrices find a complete set of orthonormal eigenvectors.
- (b) If you apply the Gram-Schmidt process (orthonormalization) to the columns of this matrix A, what is the resulting output?
- (c) If A is any m by n matrix with m > n, tell me why AA^T cannot be positive definite. Is $A^T A$ always positive definite? (If not, what is the test on A?)

7. (11 points) This problem is to find the determinants of

(a) Find $\det A$ and give a reason.

- (b) Find the cofactor C_{11} and then find det B. This is the volume of what region in \mathbb{R}^4 ?
- (c) Find $\det C$ for any value of x. You could use linearity in row 1.

8. (11 points)

- (a) When A is similar to $B = M^{-1}AM$, prove this statement: If $A^k \to 0$ when $k \to \infty$, then also $B^k \to 0$.
- (b) Suppose S is a fixed invertible 3 by 3 matrix. This question is about all the matrices A that are diagonalized by S, so that S⁻¹AS is diagonal. Show that these matrices A form a subspace of 3 by 3 matrix space. (Test the requirements for a subspace.)
- (c) Give a basis for the space of 3 by 3 diagonal matrices. Find a basis for the space in part (b)
 all the matrices A that are diagonalized by S.

9. (11 points) This square network has 4 nodes and 6 edges. On each edge, the direction of positive current $w_i > 0$ is from lower node number to higher node number. The voltages at the nodes are $(v_1, v_2, v_3, v_4.)$

- (a) Write down the incidence matrix A for this network (so that Av gives the 6 voltage differences like v_2-v_1 across the 6 edges). What is the rank of A? What is the dimension of the nullspace of A^T ?
- (b) Compute the matrix $A^T A$. What is its rank? What is its nullspace?
- (c) Suppose $v_1 = 1$ and $v_4 = 0$. If each edge contains a unit resistor, the currents $(w_1, w_2, w_3, w_4, w_5, w_6)$ on the 6 edges will be w = -Av by Ohm's Law. Then Kirchhoff's Current Law (flow in = flow out at every node) gives $A^T w = 0$ which means $A^T A v = 0$. Solve $A^T A v = 0$ for the unknown voltages v_2 and v_3 . Find all 6 currents w_1 to w_6 . How much current enters node 4?

MIT OpenCourseWare http://ocw.mit.edu

18.06 Linear Algebra Spring 2010

For information about citing these materials or our Terms of Use, visit: http://ocw.mit.edu/terms.